

PRODUCT DATA SHEET

Page 1 / 3

Lipid A from E. coli R515 (Re) TLRpure™ Sterile Solution

Cat. No.: IAX-100-004

Lot. No.:

| | |
|----------------------------|---|
| Source | Lipid A derived from E. coli R515 (Re) LPS |
| Concentration | 1mg/ml (0.5mg/ml for 250µg size) stabilised in sterile, double-distilled water (ddWater), without any additives |
| TLRpure™ | No detectable TLR4 <i>independent</i> activity: standardised potent TLR4-specific agonist |
| Purity | Ultrapure. No detectable DNA, RNA and protein traces. |
| Purification Method | R-type (mutant/rough) LPS was isolated by a phenol-chloroform-petroleum-ether method. Semi-purified LPS was subjected to further re-extraction cycles and ultracentrifugation steps, extensively electro dialysed to yield TLRpure™ LPS, from which Lipid A was generated by mild acid hydrolysis. |
| Sterility | Filter method: according to Ph. Eur. 9. Passed according to specification: <ul style="list-style-type: none"> • No growth in Thioglycolate medium at 30-35°C after 14 days. • No growth in Soybean Casein Digest Broth (TSB) at 20-25°C after 14 days. |
| Endotoxin Content | Bacterial Endotoxin Test (kinetic turbidimetric LAL method) according to Ph. Eur. 9. Endotoxin Content: >5,000,000 [EU/ml]. |
| Appearance | Colourless, clear, aqueous solution |
| Handling | Keep sterile. Prepare working dilutions from pre-warmed (~40°C) Lipid A stock solution just prior to use. Ready-to-use, sterile stock solution is cell culture-grade. No solubilisation required. Do not pre-dilute in buffer (e.g. PBS) as this will lead to precipitation of Lipid A. To yield a 100µg/ml (1,000-100x) stock solution add 100µl of Lipid A to 900µl endotoxin-free and sterile ddWater (Cat. No.: IAX-900-002) (not PBS) and mix well. |
| Activity | Optimal concentration is dependent upon cell type, species, desired activation and analysis: 0.1-1.0µg/ml <i>in vitro</i> and 5-15mg/kg <i>in vivo</i> in animal rodent models. Does not activate any TLR other than TLR4 as tested up to 1 µg/ml in relevant cellular systems (mouse macrophages). |
| Shipping | Ambient |
| Storage | 2-8°C |
| Stability | <ul style="list-style-type: none"> • Upon receipt, store product at 2-8°C. • Do not freeze. In its unopened original vial, the product is stable for at least 24 months when stored at 2-8°C. Once the glass vial is opened, or if aliquoted into sterile vials under sterile conditions, the product remains stable for an additional 12 months at 2-8°C. Pre-diluted, sterile aqueous solutions (e.g., 10-100µg/ml) are stable for a maximum of 12 hours when stored at 2-8°C due to vial surface effects on diluted solutions. |
| MSDS | Available on request |

Document No.: IAX-100-004 | **Version:** 2.1 | **Issue Date:** 01/08/2024

DISCLAIMER: THIS PRODUCT IS NOT INTENDED OR APPROVED FOR HUMAN, DIAGNOSTICS OR VETERINARY USE. USE OF THIS PRODUCT FOR HUMAN OR ANIMAL TESTING MAY BE EXTREMELY HAZARDOUS AND MAY RESULT IN DISEASE, SEVERE INJURY, OR DEATH. THIS PRODUCT IS FOR RESEARCH USE ONLY (RUO).

MATERIAL SAFETY DATA: This material should be considered hazardous until information to the contrary becomes available. Do not ingest, swallow, inhale or get into the blood stream. Do not get in eyes, on skin, or clothing. Wash thoroughly after handling. This information contains some, but not all, of the information required for the safe and proper use of this material. Access to this material must be restricted to personnel, who is appropriately experienced, qualified, competent and properly trained to use it. Material Safety Data Sheet is available upon request.

PRODUCT DATA SHEET

Page 2 / 3

Lipid A from E. coli R515 (Re) TLRpure™ Sterile Solution

Cat. No.: IAX-100-004

Lot. No.:

E. coli TLRpure™ Lipid A is a TLR4 specific agonist




FIGURE:

Macrophages from wild-type (WT) TLR4 expressing or TLR4 deficient (TLR4 KO mice were stimulated with 1 µg/ml TLRpure™ Lipid A from E. coli. Cell culture supernatants were analysed by ELISA for IL-6 after 24h. Optimal concentrations required for activation depend upon cell species (murine, human, others), cell culture conditions (FCS concentration), sampling time and cytokine analysis. Recommended range for Lipid A: 0.1-1.0 µg/ml.

Product Information

- Lipid A has been generated by mild acid hydrolysis from TLRpure™ LPS purified according to an optimised and proprietary extraction and purification protocol, but based upon the methods published by Galanos, et al. (laboratory of Westphal and Lüderitz, Freiburg, Germany).
- TLRpure™ LPS lacks any detectable bacterial, (lipo-)protein, RNA or DNA or other TLR-stimulating activity due to its ultra-purified formulation. Its unique potency and purity are quality controlled using a physiological system of primary innate immune cells and a relevant biological cytokine expression read-out.
- All immunological activity of the Lipid A is exclusively dependent upon the presence of TLR4 as determined by the use of the corresponding control cells, where TLR4 has been genetically deleted or missing (from TLR4 deficient also called TLR4 knock-out KO mice).
- TLRpure™ Lipid A convenient ready-made stabilised solution makes it the reagent of choice for in vitro as well as in vivo experiments for superior reproducible and comparable results.
- Compared to Lipid A derived from conventional (semi-purified) LPS preparations, this product is derived from low yield TLRpure™ LPS produced on an industrial fermentation scale under precisely controlled growth conditions to yield large batch sizes, allowing custom formulations/packaging.

Product Specific References

- [1] A new method for the extraction of R lipopolysaccharides. Galanos C, et al. Eur. J. Biochem. (1969); 9:245
- [2] Preparation and properties of antisera against the lipid-A component of bacterial lipopolysaccharides. Galanos C, et al. Eur. J. Biochem. (1971); 24:116
- [3] Endotoxic properties of chemically synthesized lipid A part structures. Comparison of synthetic lipid A precursor and synthetic analogues with biosynthetic lipid A precursor and free lipid A. Galanos C, et al. Eur. J. Biochem. (1984); 140:221
- [4] Lipopolysaccharides: structural principles and biologic activities. Lüderitz O, et al. Rev. Infect. Dis. (1984); 6:428
- [5] Synthetic and natural Escherichia coli free lipid A express identical endotoxic activities. Galanos C, et al. Eur. J. Biochem. (1985); 148:1

Document No.: IAX-100-004 | Version: 2.1 | Issue Date: 01/08/2024

DISCLAIMER: THIS PRODUCT IS NOT INTENDED OR APPROVED FOR HUMAN, DIAGNOSTICS OR VETERINARY USE. USE OF THIS PRODUCT FOR HUMAN OR ANIMAL TESTING MAY BE EXTREMELY HAZARDOUS AND MAY RESULT IN DISEASE, SEVERE INJURY, OR DEATH. THIS PRODUCT IS FOR RESEARCH USE ONLY (RUO).

MATERIAL SAFETY DATA: This material should be considered hazardous until information to the contrary becomes available. Do not ingest, swallow, inhale or get into the blood stream. Do not get in eyes, on skin, or clothing. Wash thoroughly after handling. This information contains some, but not all, of the information required for the safe and proper use of this material. Access to this material must be restricted to personnel, who is appropriately experienced, qualified, competent and properly trained to use it. Material Safety Data Sheet is available upon request.

PRODUCT DATA SHEET

Page 3 / 3

Lipid A from E. coli R515 (Re) TLRpure™ Sterile Solution

Cat. No.: IAX-100-004

Lot. No.:

General Information

- Activation of cells by LPS is mediated by the Toll-like receptor 4 (TLR4), a member of the highly conserved protein family of TLRs, which are specialised in the recognition of microbial components. In mice, defects in TLR4 result in LPS unresponsiveness.
- For optimal interaction with LPS, TLR4 requires association with myeloid differentiation protein 2 (MD-2). According to current consensus activation of TLR4 is preceded by the transfer of LPS to membrane-bound (m) or soluble (s) CD14 by LPS-binding protein (LBP). This mechanism is believed to be generally true for LPS signaling. Re-form LPS and lipid A, but not S-form LPS, are capable of inducing TNF- α responses also in the absence of CD14.
- LPS, synthesised by most wild-type (WT) Gram-negative bacteria (S-form LPS), consists of three regions, the O-polysaccharide chain, which is made up of repeating oligosaccharide units, the core oligosaccharide and the lipid A, which harbors the endotoxic activity of the entire molecule. R-form LPS synthesised by the so-called rough (R) mutants of Gram-negative bacteria lacks the O-specific chain. Furthermore, the core-oligosaccharide may be present in different degrees of completion, depending on the class (Ra to Re) to which the mutant belongs. Monophosphoryl Lipid A (MPLA) represents a detoxified derivative of Lipid A and constitutes an important adjuvant in prophylactic and therapeutic vaccines.

References

- [1] *R-form LPS, the master key to the activation of TLR4/MD-2-positive cells.* Huber M, et al. Eur. J. Immunol. (2006); 36:701
- [2] *CD14 is required for MyD88-independent LPS signaling.* Jiang Z, Georgel P, Du X, Shamel L, Sovath S, Mudd S, Huber M, Kalis C, Keck S, Galanos C, Freudenberg M, Beutler B. Nat. Immunol. (2005); 6:565
- [3] *Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: mutations in Tlr4 gene.* Poltorak A, He X, Smirnova I, Liu MY, Van Huffel C, Du X, Birdwell D, Alejos E, Silva M, Galanos C, Freudenberg M, Ricciardi-Castagnoli P, Layton B, Beutler B. Science (1998); 282:2085
- [4] *Structural relationship of Salmonella O and R antigens.* Lüderitz O, Galanos C, et al. Ann. N.Y. Acad. Sci. (1966); 133:349
- [5] *Preparation and properties of antisera against the lipid-A component of bacterial lipopolysaccharides.* Galanos C, et al. Eur. J. Biochem. (1971); 24:116
- [6] *Lipid A: chemical structure and biological activity.* Lüderitz O, Galanos C, et al. J. Infect. Dis. (1973); 128:17
- [7] *Purification and structural determination of nontoxic lipid A obtained from the lipopolysaccharide of Salmonella typhimurium.* Qureshi N, Takayama K, Ribí E. J. Biol. Chem. (1982); 257:11808

Document No.: IAX-100-004 | **Version:** 2.1 | **Issue Date:** 01/08/2024

DISCLAIMER: THIS PRODUCT IS NOT INTENDED OR APPROVED FOR HUMAN, DIAGNOSTICS OR VETERINARY USE. USE OF THIS PRODUCT FOR HUMAN OR ANIMAL TESTING MAY BE EXTREMELY HAZARDOUS AND MAY RESULT IN DISEASE, SEVERE INJURY, OR DEATH. THIS PRODUCT IS FOR RESEARCH USE ONLY (RUO).

MATERIAL SAFETY DATA: This material should be considered hazardous until information to the contrary becomes available. Do not ingest, swallow, inhale or get into the blood stream. Do not get in eyes, on skin, or clothing. Wash thoroughly after handling. This information contains some, but not all, of the information required for the safe and proper use of this material. Access to this material must be restricted to personnel, who is appropriately experienced, qualified, competent and properly trained to use it. Material Safety Data Sheet is available upon request.